

VU Research Portal

Developmental Origins of Increased Nuchal Translucency

Burger, N.B.

2016

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Burger, N. B. (2016). *Developmental Origins of Increased Nuchal Translucency*. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam].

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Chapter 1

General introduction and outline of the thesis



INTRODUCTION

The nuchal translucency (NT) is a subcutaneous collection of fluid in the posterior neck region of a fetus¹. It can be visualized by ultrasound at 10-14 weeks of gestation and disappears in the majority of the fetuses after 14 weeks of gestation². An increased NT is defined as a measurement of the translucent area in the neck region above the 95th percentile for the gestational age³. Ultrasound measurement of NT is an important part of first-trimester screening for trisomy 21, trisomy 18 and trisomy 13^{4,5}. Increased NT is also associated with monosomy X⁶. In euploid fetuses an increased NT identifies fetuses at risk for cardiac defects⁷⁻¹³, other structural malformations^{10,12}, rare genetic syndromes and skeletal dysplasias¹⁰⁻¹². Nevertheless, the majority of euploid fetuses with increased NT have a healthy, uncomplicated outcome¹⁴.

Although NT measurement is used as part of first-trimester screening, a pathophysiological explanation for increased NT is lacking. Several mechanisms have been suggested, but they fail to explain both the regional and temporary character of nuchal edema, as well as the wide spectrum of fetal anomalies associated with increased NT.

Initially, it was believed that increased NT originates from cardiac failure, either due to myocardial dysfunction or secondary to a cardiac defect¹⁵. This hypothesis has been based on the high frequency of cardiac defects in euploid and aneuploid fetuses with increased NT^{7-13,16-21} and the fact that the risk of a cardiac defect increases with enlargement in NT in euploid fetuses¹⁷. But not all fetuses with increased NT have a cardiac defect. Furthermore, the most common cardiac defect in fetuses with increased NT is a ventricular septal defect. This type of cardiac defect does not cause cardiac failure in fetal life. In fact, the majority of cardiac defects do not compromise fetal hemodynamics²². If cardiac failure causes increased NT, then specific types of cardiac defects that can result in fetal hemodynamic alterations – such as tricuspid valve insufficiency or stenosis – should be overrepresented in fetuses with increased NT. Enlarged NT is, however, not related to a specific type of cardiac defect, but is related to any type of cardiac defect^{17,22-24}. Besides, fetal cardiac failure is usually progressive towards fetal death and is not a temporary phenomenon.

Cardiac failure has also been suggested to cause increased NT because of the finding of abnormal ductus venosus flow velocity waveforms in fetuses with increased NT^{15,25-28}. It was suggested that cardiac failure causes both increased NT and abnormal ductal flow velocity waveforms. But intra-cardiac flow velocities do not differ between euploid and aneuploid fetuses with normal or enlarged NT, irrespective of cardiac anatomy²⁹. The fact that abnormal ductal flow velocity waveforms are also observed in euploid fetuses with normal NT, showing both normal and abnormal cardiac anatomy^{30,31}, and in aneuploid fetuses with increased NT and normal cardiac anatomy³² do not support the theory of cardiac failure as an explanation for increased NT.

If cardiac failure causes increased NT, then fetal findings of cardiac decompensation, such as pleural and pericardial effusion, ascites and cardiomegaly, would be expected. These findings are, however, usually not observed in fetuses with increased NT. Cardiac failure can neither

explain the local and temporary fluid accumulation in the neck region. In conclusion, the contention of cardiac failure as explanation for increased NT is commonly presumed but is based on assumptions and is supported by neither evidence nor pathophysiology.

Previous studies showed abnormally enlarged jugular lymphatic sacs in the neck of fetuses with increased NT, together with the finding that increased NT is in fact mesenchymal edema³³⁻³⁵. It was suggested that a disturbance in lymphatic development caused an increased NT. A longitudinal ultrasound study in fetuses with increased NT showed a temporal relationship between the development of NT and the volume of the enlarged jugular lymphatic sacs, in which increased NT preceded the jugular lymphatic sac expansion³⁶. Lymphatic development starts in the neck by the formation of the jugular lymphatic sacs³⁷. Lymphatic endothelial cells bud and migrate from the cardinal veins as single cells or clusters and subsequently form the jugular lymphatic sacs³⁸. The jugular lymphatic sacs then remodel into lymph nodes after 10 weeks of gestation³⁹. The development of the lymphatic system is completed by the ingrowth of the right thoracic duct into the left jugular lymphatic sac, thereby forming the main drainage site of lymphatic fluid into the venous systemic circulation^{39,40}. A delay or disturbance in lymphatic development could explain the temporary and local fluid accumulation in the nuchal region. Immunohistochemical studies showed a diminished expression of lymphatic markers Prox-1 and Podoplanin and an increased expression of blood vessel markers, such as Neuropilin-1, Smooth Muscle Actin (SMA) and Vascular Endothelial Growth Factor (VEGF)-A in the jugular lymphatic sacs in aneuploid mouse embryos and human fetuses with nuchal edema⁴¹⁻⁴³. Abnormal (lymphatic) endothelial development could explain the finding of enlarged jugular lymphatic sacs and nuchal edema. Morphological studies in trisomy 16 mouse embryos with nuchal edema demonstrated an abnormal development and altered positioning of cranial nerves, which are consistently located adjacent to the jugular lymphatic sacs⁴⁴. It was suggested that abnormal neuronal development could disturb lymphatic endothelial differentiation of the jugular lymphatic sacs⁴⁴. Cranial nerves are derived from the neural crest and develop in the same time period in the late first trimester as nuchal edema⁴⁵. Disturbances in neural crest cell migration or differentiation cause cardiac defects⁴⁶⁻⁴⁸, craniofacial abnormalities⁴⁷ and skeletal malformations⁴⁹, which are anomalies that are often found in fetuses with increased NT.

In summary, a lymphatic developmental disorder seems to be involved in the etiology of increased NT. Disturbed endothelial and neuronal development seem to be prominent denominators in the pathophysiology of increased NT.

AIM AND OUTLINE OF THE THESIS

This thesis describes the translational approach to explain the clinical finding of fetal increased NT. The aim of this thesis was to study the relationship between increased NT, lymphatic development, cardiovascular development and the involvement of nerves to provide more insight into the developmental background of increased NT.

In **chapter 2** we have identified genes involved in normal cardiac and lymphatic development. Loss of function of these genes causes both cardiac and lymphatic abnormalities in mouse embryos with nuchal edema. This provides insight into the underlying genetic mechanisms that can explain the high incidence of cardiac defects in fetuses with increased NT.

In **chapter 3** the involvement of cranial nerve progenitor cells, the neural crest cells, has been investigated in lymphatic development. Retinoic acid is synthesized in nerves and is essential for neural crest cell and lymphatic development. The involvement of retinoic acid at specific stages of lymphatic differentiation and in nuchal edema has been examined.

In **chapter 4** the theory of cardiac failure as explanation for increased NT has been investigated. The nuchal region, jugular lymphatic sacs and cardiac anatomy have been studied in various mutant mouse models with lymphatic and cardiac defects with and without the presence of nuchal edema.

Altered hemodynamics, reflected by abnormal ductus venosus flow velocity waveforms, has been suggested to be involved in the development of increased NT. **Chapter 5** describes the use of a uniform anatomical definition for the ductus venosus. In **chapter 6** the systematic morphological development of the ductus venosus and the presence of a ductus venosus sphincter have been studied in wild-type mouse and human embryos. In **chapter 7** we have investigated whether a local alteration in ductus venosus morphology causes abnormal ductus venosus flow velocity waveforms in several mutant mouse models with lymphatic and cardiac defects with and without the presence of nuchal edema. In **chapter 8** the origin of abnormal ductus venosus flow velocity waveforms and its relationship with increased NT and cardiac defects has been further explored in human fetuses. Postmortem examination of the ductus venosus, heart and jugular lymphatic sacs has been performed in human fetuses with normal and abnormal ductus venosus flow velocity waveforms, showing normal and increased NT. This thesis ends with a general discussion and future perspectives in **chapter 9**. A summary of the presented data is given in both English and Dutch in **chapter 10**.

REFERENCES

- Nicolaides KH, Azar G, Byrne D, Mansur C, Marks K. Fetal nuchal translucency: Ultrasound screening for chromosomal defects in first trimester of pregnancy. *BMJ* 1992;304:867-869.
- Pajkrt E, De Graaf IM, Mol BWJ, Van Lith JMM, Bleker OP, Bilardo CM. Weekly nuchal translucency measurements in normal fetuses. *Obstet Gynecol* 1998;91:208-211.
- Pajkrt E, Bilardo CM, Van Lith JMM, Mol BWJ, Bleker OP. Nuchal translucency measurement in normal fetuses. *Obstet Gynecol* 1995;86:994-997.
- Pandya PP, Brizot ML, Kuhn P, Snijders RJM, Nicolaides KH. First-trimester fetal nuchal translucency thickness and risk for trisomies. *Obstet Gynecol* 1994;84:420-423.
- Snijders RJ, Noble P, Sebire N, Souka A, Nicolaides KH. UK multicentre project on assessment of risk of trisomy 21 by maternal age and fetal nuchal-translucency thickness at 10-14 weeks of gestation. *Lancet* 1998;352:343-346.
- Pandya PP, Kondylis A, Hilbert L, Snijders RJ, Nicolaides KH. Chromosomal defects and outcome in 1015 fetuses with increased nuchal translucency. *Ultrasound Obstet Gynecol* 1995;5:15-19.
- Hyett J, Moscoso G, Papapanagiotou G, Perdu M, Nicolaides KH. Abnormalities of the heart and great arteries in chromosomally normal fetuses with increased nuchal translucency thickness at 11-13 weeks of gestation. *Ultrasound Obstet Gynecol* 1996;7:245-250.
- Hyett JA, Perdu M, Sharland GK, Snijders RSM, Nicolaides KH. Increased nuchal translucency at 10-14 weeks of gestation as a marker for major cardiac defects. *Ultrasound Obstet Gynecol* 1997;10:242-246.
- Hafner E, Schuchter K, Liebhart E, Philipp K. Results of routine fetal nuchal translucency measurement at weeks 10-13 in 4233 unselected pregnant women. *Prenat Diagn* 1998;18:29-34.
- Souka AP, Snijders RJM, Novakov A, Soares W, Nicolaides KH. Defects and syndromes in chromosomally normal fetuses with increased nuchal translucency thickness at 10-14 weeks of gestation. *Ultrasound Obstet Gynecol* 1998;11:391-400.
- Souka AP, von Kaisenberg CS, Hyett JA, Sonek JD, Nicolaides KH. Increased nuchal translucency with normal karyotype. *Am J Obstet Gynecol* 2005;192:1005-1021.
- Souka AP, Krampl E, Bakalis S, Heath V, Nicolaides KH. Outcome of pregnancy in chromosomally normal fetuses with increased nuchal translucency in the first trimester. *Ultrasound Obstet Gynecol* 2001;18:9-17.
- Westin M, Saltvedt S, Bergman G, Almstrom H, Grunewald C, Valentin L. Is measurement of nuchal translucency thickness a useful screening tool for heart defects? A study of 16,383 fetuses. *Ultrasound Obstet Gynecol* 2006;27:632-639.
- Van Vugt JMG, Tinnemans BW, Zalen-Sprock RMv. Outcome and early childhood follow-up of chromosomally normal fetuses with increased nuchal translucency at 10-14 weeks' gestation. *Ultrasound Obstet Gynecol* 1998;11:407-409.
- Montenegro N, Matias A, Areias JC, Castedo S, Barros H. Increased fetal nuchal translucency: Possible involvement of early cardiac failure. *Ultrasound Obstet Gynecol* 1997;10:265-268.
- Bilardo CM, Pajkrt E, De Graaf IM, Mol BW, Bleker OP. Outcome of fetuses with enlarged nuchal translucency and normal karyotype. *Ultrasound Obstet Gynecol* 1998;11:401-406.
- Ghi T, Huggon IC, Zosmer N, Nicolaides KH. Incidence of major structural cardiac defects associated with increased nuchal translucency but normal karyotype. *Ultrasound Obstet Gynecol* 2001;18:610-614.
- Hyett JA, Moscoso G, Nicolaides KH. First-trimester nuchal translucency and cardiac septal defects in fetuses with trisomy 21. *Am J Obstet Gynecol* 1995;172:1411-1413.
- Hyett J, Moscoso G, Nicolaides K. Abnormalities of the heart and great arteries in first trimester chromosomally abnormal fetuses. *Am J Med Genet* 1997;69:207-216.
- Hyett JA, Moscoso G, Nicolaides KH. Cardiac defects in 1st-trimester fetuses with trisomy 18. *Fetal Diagn Ther* 1995;10:381-386.
- Vogel M, Sharland GK, McElhinney DB, Zidere V, Simpson JM, Miller OI, Allan LD. Prevalence of increased nuchal translucency in fetuses with congenital cardiac disease and a normal karyotype. *Cardiol Young* 2009;19:441-445.
- Simpson JM, Sharland GK. Nuchal translucency and congenital heart defects: heart failure or not? *Ultrasound Obstet Gynecol* 2000;16:30-36.
- de Mooij YM, Haak MC, Bartelings MM, Twisk JW, Gittenberger-de GA, van Vugt JM, Bekker MN. Abnormal ductus venosus flow in first-trimester fetuses with increased nuchal translucency: relationship with the type of cardiac defect? *J Ultrasound Med* 2010;29:1051-1058.
- Clur SA, Mathijssen IB, Pajkrt E, Cook A, Laurini RN, Ottenkamp J, Bilardo CM. Structural heart defects associated with an increased nuchal translucency: 9 years experience in a referral centre. *Prenat Diagn* 2008;28:347-354.
- Matias A, Gomes C, Flack N, Montenegro N, Nicolaides KH. Screening for chromosomal abnormalities at 10-14 weeks: the role of ductus venosus blood flow. *Ultrasound Obstet Gynecol* 1998;12:380-384.
- Borrell A, Antolin E, Costa D, Farre MT, Martinez JM, Fortuny A. Abnormal ductus venosus blood flow in trisomy 21 fetuses during early pregnancy. *Am J Obstet Gynecol* 1998;179:1612-1617.
- Bilardo CM, Muller MA, Zikulnig L, Schipper M, Hecher K. Ductus venosus studies in fetuses at high risk for chromosomal or heart abnormalities: relationship with nuchal translucency measurement and fetal outcome. *Ultrasound Obstet Gynecol* 2001;17:288-294.
- Matias A, Huggon I, Areias JC, Montenegro N, Nicolaides KH. Cardiac defects in chromosomally normal fetuses with abnormal ductus venosus blood flow at 10-14 weeks. *Ultrasound Obstet Gynecol* 1999;14:307-310.
- Haak MC, Twisk JWR, Bartelings MM, Gittenberger-de Groot, Van Vugt JMG. First-trimester fetuses with increased nuchal translucency do not show changed intracardiac flow velocities. *Ultrasound Obstet Gynecol* 2005;25:246-52.
- Oh C, Harman C, Baschat AA. Abnormal first-trimester ductus venosus blood flow: a risk factor for adverse outcome in fetuses with normal nuchal translucency. *Ultrasound Obstet Gynecol* 2007;30:192-196.
- Martinez JM, Comas M, Borrell A, Bannasar M, Gomez O, Puerto B, Gratacos E. Abnormal first-trimester ductus venosus blood flow: a marker of cardiac defects in fetuses with normal karyotype and nuchal translucency. *Ultrasound Obstet Gynecol* 2010;35:267-272.
- Haak MC, Twisk JW, Bartelings MM, Gittenberger-de Groot, van Vugt JM. Ductus venosus flow velocities in relation to the cardiac defects in first-trimester fetuses with enlarged nuchal translucency. *Am J Obstet Gynecol* 2003;188:727-733.
- Haak MC, Bartelings MM, Jackson DG, Webb S, Van Vugt JMG, Gittenberger-de Groot. Increased nuchal translucency is associated with jugular lymphatic distension. *Hum Reprod* 2002;17:1086-1092.
- Bekker MN, Haak MC, Rekoert-Hollander M, Twisk JWR, Van Vugt JMG. Increased nuchal translucency and distended jugular lymphatic sacs by first-trimester ultrasound. *Ultrasound Obstet Gynecol* 2005;25:239-45.
- Gittenberger-de Groot AC, Van Den Akker NM, Bartelings MM, Webb S, Van Vugt JMG, Haak MC. Abnormal lymphatic development in trisomy 16 mouse embryos precedes nuchal edema. *Dev Dyn* 2004;230:378-384.
- Bekker MN, Twisk JW, Bartelings MM, Gittenberger-de Groot, van Vugt JM. Temporal relationship between increased nuchal translucency and enlarged jugular lymphatic sac. *Obstet Gynecol* 2006;108:846-853.
- Srinivasan RS, Dillard ME, Lagutin OV, Lin FJ, Tsai S, Tsai MJ, Samokhvalov IM, Oliver G. Lineage tracing demonstrates the venous origin of the mammalian lymphatic vasculature. *Genes Dev* 2007;21:2422-2432.
- Hagerling R, Pollmann C, Andreas M, Schmidt C, Nurmi H, Adams RH, Alitalo K, Andresen V, Schulte-Merker S, Kiefer F. A novel multistep mechanism for initial lymphangiogenesis in mouse embryos based on ultramicroscopy. *EMBO J* 2013;32:629-644.
- Sabin FR. The lymphatic system in human embryos, with a consideration of the morphology of the system as a whole. *Am J Anat* 1909;9:43-91.
- Van der Putte SC. The development of the lymphatic system in man. *Adv Anat Embryol Cell Biol* 1975;51:3-60.
- Bekker MN, Van Den Akker NM, Bartelings MM, Arksteijn JB, Fischer SG, Polman JA, Haak MC, Webb S, Poelmann RE, van Vugt JM, Gittenberger-de Groot. Nuchal edema and venous-lymphatic phenotype disturbance in human fetuses and mouse embryos with aneuploidy. *J Soc Gynecol Investig* 2006;13:209-216.
- de Mooij YM, Van Den Akker NM, Bekker MN, Bartelings MM, Wisse LJ, van Vugt JM, Gittenberger-de Groot. Abnormal Shh and FOXC2 expression correlates with aberrant lymphatic development in human fetuses with increased nuchal translucency. *Prenat Diagn* 2009;29:840-846.
- de Mooij YM, Van Den Akker NM, Bekker MN, Bartelings MM, van Vugt JM, Gittenberger-de Groot. Aberrant lymphatic development in euploid fetuses with increased nuchal translucency including Noonan syndrome. *Prenat Diagn* 2011;31:159-166.
- Bekker MN, Arksteijn JB, Van Den Akker NM, Moulin DGM, Hoffman S, Webb S, Van Vugt JMG, Gittenberger-de Groot. Increased NCAM expression and vascular development in trisomy 16 mouse embryos: relationship with nuchal translucency. *Pediatr Res* 2005;58:1222-7.

45. Muller F, O'Rahilly R. The initial appearance of the cranial nerves and related neuronal migration in staged human embryos. *Cells Tissues Organs* 2011;193:215-238.46. Jiang X, Rowitch DH, Soriano P, McMahon AP, Sucov HM. Fate of the mammalian cardiac neural crest. *Development* 2000;127:1607-1616.
47. Huang ZP, Chen JF, Regan JN, Maguire CT, Tang RH, Dong XR, Majesky MW, Wang DZ. Loss of microRNAs in neural crest leads to cardiovascular syndromes resembling human congenital heart defects. *Arterioscler Thromb Vasc Biol* 2010;30:2575-2586.
48. Kirby ML, Waldo KL. Neural crest and cardiovascular patterning. *Circ Res* 1995;77:211-215.
49. Van Ho AT, Hayashi S, Brohl D, Aurade F, Rattenbach R, Relaix F. Neural crest cell lineage restricts skeletal muscle progenitor cell differentiation through Neuregulin1-ErbB3 signaling. *Dev Cell* 2011;21:273-287.